Ohio Primary and Secondary Syphilis Cases, 2010 to First Quarter 2012
Demographic Breakdown of Ohio
Presented by Date of Diagnosis[1]

	Cases Reported to the Ohio Department of Health								
	Jan - March Cases			January - December Cases				Rate[2]	
AGE	**2010**	**2011**	**2012**	**2010**	**% of Total**	**2011**	**% of Total**	**2010**	**2011**
0 - 9	0	0	0	0	0.0	0	0.0	0.0	0.0
10 - 14	1	1	0	3	0.6	2	0.5	0.4	0.3
15 - 19	16	10	8	69	13.0	57	12.9	8.4	6.9
20 - 24	31	35	20	132	25.0	126	28.4	17.3	16.5
25 - 29	16	22	16	99	18.7	80	18.1	13.8	11.1
30 - 34	16	19	11	73	13.8	53	12.0	10.6	7.7
35 - 39	4	6	7	38	7.2	33	7.4	5.3	4.6
40 - 44	11	9	5	41	7.8	37	8.4	5.4	4.9
45 - 54	13	12	16	59	11.2	39	8.8	3.4	2.2
55 - 64	2	4	3	15	2.8	13	2.9	1.0	0.9
65+	0	0	1	0	0.0	3	0.7	0.0	0.2
Not Specified	0	0	0	0	0.0	0	0.0	--	--
RACE									
Am. Indian/Alaskan Native	0	0	0	1	0.2	1	0.2	2.6	2.6
Asian/Pacific Islander	0	0	0	1	0.2	0	0.0	0.5	0.0
Black/African American	71	72	49	352	66.5	292	65.9	23.3	19.3
White	36	44	34	167	31.6	135	30.5	1.7	1.4
Other[3]	3	2	4	7	1.3	13	2.9	--	--
Not Specified	0	0	0	1	0.2	2	0.5	--	--
ETHNICITY									
Hispanic	1	3	5	8	1.5	6	1.4	2.3	1.7
Non-Hispanic	109	115	82	519	98.1	435	98.2	4.6	3.9
Not Specified	0	0	0	2	0.4	2	0.5	--	--
SEX									
Male	82	95	70	398	75.2	335	75.6	6.7	5.7
Female	28	23	17	131	24.8	108	24.4	--	--
Not Specified	0	0	0	0	0.0	0	0.0	--	--
TOTAL	**110**	**118**	**87**	**529**	**100.0**	**443**	**100.0**	**4.6**	**3.8**

Notes:
(1) In this presentation, date of diagnosis is defined as the date the specimen was collected by the provider.
(2) Rates are shown per 100,000 persons and were calculated using census estimates for 2010.
(3) Cases are included in the Other race category if multiple races are chosen or if the case did not fit into any of the listed categories.
Small numbers are unstable and should be interpreted with caution.
Provisional data. Numbers subject to change when additional information is gained.

Source: Ohio Department of Health, STD Surveillance. Data reported through 06/10/12.

Ohio Primary and Secondary Syphilis Cases, 2010 to First Quarter 2012
Presented by Date of Diagnosis[1]

COUNTY	January - March Cases 2010	2011	2012	Jan - Dec Cases 2010	2011	Rate[2] 2010	2011
ADAMS	0	0	0	0	1	0.0	3.5
ALLEN	0	0	0	0	0	0.0	0.0
ASHLAND	0	0	0	0	0	0.0	0.0
ASHTABULA	0	0	0	0	0	0.0	0.0
ATHENS	0	0	0	0	0	0.0	0.0
AUGLAIZE	0	0	0	0	0	0.0	0.0
BELMONT	0	0	0	0	1	0.0	1.4
BROWN	0	0	0	1	0	2.2	0.0
BUTLER	4	4	3	9	9	2.4	2.4
CARROLL	0	0	0	1	0	3.5	0.0
CHAMPAIGN	0	0	0	0	0	0.0	0.0
CLARK	1	0	0	2	1	1.4	0.7
CLERMONT	0	1	2	1	2	0.5	1.0
CLINTON	0	0	0	0	0	0.0	0.0
COLUMBIANA	0	0	0	0	0	0.0	0.0
COSHOCTON	0	0	0	3	0	8.1	0.0
CRAWFORD	0	0	0	1	0	2.3	0.0
CUYAHOGA	6	14	12	70	53	5.5	4.1
DARKE	0	0	0	0	1	0.0	1.9
DEFIANCE	0	0	0	0	0	0.0	0.0
DELAWARE	0	0	0	2	1	1.1	0.6
ERIE	0	0	0	1	1	1.3	1.3
FAIRFIELD	1	0	0	4	0	2.7	0.0
FAYETTE	0	0	0	0	0	0.0	0.0
FRANKLIN	24	44	25	109	116	9.4	10.0
FULTON	0	0	0	0	0	0.0	0.0
GALLIA	0	0	0	0	0	0.0	0.0
GEAUGA	0	0	0	1	0	1.1	0.0
GREENE	0	0	0	1	2	0.6	1.2
GUERNSEY	0	0	0	0	0	0.0	0.0
HAMILTON	54	43	31	245	195	30.5	24.3
HANCOCK	0	0	0	0	1	0.0	1.3
HARDIN	0	0	0	0	0	0.0	0.0
HARRISON	0	0	0	0	0	0.0	0.0
HENRY	0	0	0	0	0	0.0	0.0
HIGHLAND	0	0	0	0	0	0.0	0.0
HOCKING	0	0	0	2	0	6.8	0.0
HOLMES	0	0	1	0	0	0.0	0.0
HURON	0	0	0	0	0	0.0	0.0
JACKSON	1	0	0	1	0	3.0	0.0
JEFFERSON	0	0	0	1	0	1.4	0.0
KNOX	0	0	0	0	0	0.0	0.0
LAKE	1	0	0	4	0	1.7	0.0
LAWRENCE	0	0	0	0	0	0.0	0.0
LICKING	0	2	1	3	5	1.8	3.0

Notes:

cont.

(1) In this presentation, date of diagnosis is defined as the date the specimen was collected by the provider.

(2) Rates are shown per 100,000 persons and were calculated using census estimates for 2010.

Small numbers are unstable and should be interpreted with caution.

Provisional data. Numbers subject to change when additional information is gained.

Source: Ohio Department of Health, STD Surveillance. Data reported through 06/10/12.

Ohio Primary and Secondary Syphilis Cases, 2010 to First Quarter 2012
Presented by Date of Diagnosis[1]

COUNTY	January - March Cases 2010	2011	2012	Jan - Dec Cases 2010	2011	Rate[2] 2010	2011
LOGAN	0	0	0	1	0	2.2	0.0
LORAIN	0	1	0	8	2	2.7	0.7
LUCAS	2	0	3	9	9	2.0	2.0
MADISON	0	0	1	1	0	2.3	0.0
MAHONING	2	1	1	3	3	1.3	1.3
MARION	0	0	0	2	0	3.0	0.0
MEDINA	0	1	0	0	1	0.0	0.6
MEIGS	0	0	0	0	0	0.0	0.0
MERCER	0	0	0	0	0	0.0	0.0
MIAMI	0	0	0	0	0	0.0	0.0
MONROE	0	0	0	0	0	0.0	0.0
MONTGOMERY	3	2	5	8	17	1.5	3.2
MORGAN	0	0	0	0	0	0.0	0.0
MORROW	0	0	0	0	0	0.0	0.0
MUSKINGUM	0	0	0	1	0	1.2	0.0
NOBLE	0	0	0	1	0	6.8	0.0
OTTAWA	0	0	0	0	0	0.0	0.0
PAULDING	0	0	0	0	0	0.0	0.0
PERRY	0	0	0	1	0	2.8	0.0
PICKAWAY	0	0	0	0	0	0.0	0.0
PIKE	0	0	0	0	0	0.0	0.0
PORTAGE	0	0	0	0	3	0.0	1.9
PREBLE	0	0	0	0	0	0.0	0.0
PUTNAM	0	0	0	0	0	0.0	0.0
RICHLAND	2	1	0	2	1	1.6	0.8
ROSS	1	0	0	1	0	1.3	0.0
SANDUSKY	0	0	0	3	0	4.9	0.0
SCIOTO	1	0	0	3	1	3.8	1.3
SENECA	0	0	0	0	0	0.0	0.0
SHELBY	0	0	1	0	0	0.0	0.0
STARK	0	1	0	3	4	0.8	1.1
SUMMIT	5	2	0	16	7	3.0	1.3
TRUMBULL	0	0	0	1	0	0.5	0.0
TUSCARAWAS	1	0	0	1	0	1.1	0.0
UNION	0	0	0	0	1	0.0	1.9
VAN WERT	0	0	0	0	0	0.0	0.0
VINTON	0	0	0	0	0	0.0	0.0
WARREN	0	1	1	0	3	0.0	1.4
WASHINGTON	0	0	0	1	0	1.6	0.0
WAYNE	0	0	0	0	1	0.0	0.9
WILLIAMS	0	0	0	0	0	0.0	0.0
WOOD	1	0	0	1	0	0.8	0.0
WYANDOT	0	0	0	0	0	0.0	0.0
UNKNOWN COUNTY[3]	0	0	0	0	1	--	--
TOTAL	**110**	**118**	**87**	**529**	**443**	**4.6**	**3.8**

Notes:
(1) In this presentation, date of diagnosis is defined as the date the specimen was collected by the provider.
(2) Rates are shown per 100,000 persons and were calculated using census estimates for 2010.
(3) Cases reported without patient's county are classified as "Unknown County".
Small numbers are unstable and should be interpreted with caution.
Provisional data. Numbers subject to change when additional information is gained.

Source: Ohio Department of Health, STD Surveillance. Data reported through 06/10/12.

Ohio Primary and Secondary Syphilis Cases, 2010 to First Quarter 2012
Presented by Date of Diagnosis[1]

CITY	January - March Cases 2010	2011	2012	Jan - Dec Cases 2010	2011	Rate[2] 2010	2011
Akron	4	2	0	11	6	5.5	3.0
Canton	0	1	0	1	3	1.4	4.1
Cincinnati	54	42	31	241	192	81.2	64.7
Cleveland	6	10	7	54	37	13.6	9.3
Columbus	20	38	23	99	104	12.9	13.5
Dayton	2	2	4	6	17	4.2	12.0
Lima	0	0	0	7	9	18.1	23.2
Toledo	2	0	2	7	9	2.4	3.1
Youngstown	2	1	0	3	2	4.5	3.0



Notes:
(1) In this presentation, date of diagnosis is defined as the date the specimen was collected by the provider.
(2) Rates are shown per 100,000 persons and were calculated using census estimates for 2010.
Small numbers are unstable and should be interpreted with caution.
Provisional data. Numbers subject to change when additional information is gained.

Source: Ohio Department of Health, STD Surveillance. Data reported through 06/10/12.